

Claims

1. A method to prepare a desired polyketide synthase from individual modules which method comprises  
providing successive covalently linked modules that comprise an intrapeptide  
linker (RAL) and successively non-covalently linked modules that comprise  
interpeptide linkers (ERL), so as to facilitate the transfer of a nascent polyketide  
chain from the a module of said PKS to a succeeding module of said PKS; and  
assembling said modules.

2. The method of claim 1 wherein said assembling is by incubating the  
polyketides which comprise said modules in a reaction mixture.

3. The method of claim 1 wherein said assembling is by expressing  
constructs which encode said modules in a host cell.

4. The method of claim 1 wherein at least some of the modules in the desired  
polyketide synthase are derived from a library of Type I PKS modules and wherein at  
least one module in said polyketide synthase is heterologous with respect to the  
remaining modules.

5. The method of claim 1 wherein each RAL has an amino acid sequence  
selected from the group consisting of those set forth in Figure 3 or a variant thereof and  
the N-terminal portions of each ERL has the amino acid sequence set forth in Figure 3 or  
a variant thereof.

6. A polyketide synthase prepared by the method of claim 4.

7. The polyketide synthase of claim 6 which contains erythromycin modules 1, and 3-6 and tylosin module 2, and wherein said polyketide chain is transferred from *ery* module 1 to *tyl* module 2 and then to *ery* modules 3-6.

5 8. The polyketide synthase of claim 6 which contains erythromycin modules 1-5 and narbomycin module 6, wherein said polyketide chain is passed from *ery* modules 1-5 to *nar* module 6.

10 9. The polyketide synthase of claim 6 which contains modules 1 and 3-6 of erythromycin and modules 2-3 of tylosin, spiramycin or niddamycin, wherein said polyketide chain is passed from *ery* module 1 to modules 2-3 of tylosin, spiramycin or niddamycin and then to *ery* modules 3-6.

15 10. The polyketide synthase of claim 6 which contains modules 1-3 of tylosin, spiramycin or niddamycin and modules 3-6 of erythromycin, and wherein said polyketide chain is passed from modules 1-3 of said tylosin, spiramycin or niddamycin to *ery* modules 3-6.

20 11. The polyketide synthase of claim 6 which contains a module of tylosin, spiramycin or niddamycin and modules 1-2 and 3-6 of erythromycin, wherein said polyketide chain is passed from *ery* modules 1-2 to the tylosin, spiramycin or niddamycin module and then to *ery* modules 3-6.

25 12. The polyketide synthase of claim 6 which contains modules 1 and 3-6 of erythromycin and module 5 of tylosin, spiramycin or niddamycin having the enoyl reductase catalytic activity inactivated, wherein said polyketide chain is passed from *ery* module 1 to module 5 of tylosin, spiramycin or niddamycin and then to *ery* modules 3-6.

30 13. The polyketide synthase of claim 6 which contains erythromycin modules 1-4 and 6 and module 6 of spiramycin or niddamycin, wherein said polyketide

chain is passed from *ery* modules 1-4 to module 6 of spiramycin or niddamycin and then to *ery* module 6.

14. The polyketide synthase of claim 6 which contains module 1 of  
FK-506/520 and modules 2-14 of rapamycin, wherein said polyketide chain is passed  
from module 1 of FK-506/520 and then to modules 2-14 of rapamycin.

15. The polyketide synthase of claim 6 which contains module 1 and 11-14 of  
rapamycin and modules 2-6 of FK-506/520 wherein said polyketide chain is passed from  
module 1 of rapamycin to modules 2-6 of FK-506/520 and then to modules 11-14 of  
rapamycin.

16. The polyketide synthase of claim 6 which contains module 1 of  
rapamycin, modules 2-7 of FK-506/520 and modules 12-14 of rapamycin, wherein said  
polyketide chain is passed from module 1 of rapamycin to modules 2-7 of FK-506/520  
and then to modules 12-14 of rapamycin.

17. The polyketide synthase of claim 6 which contains module 1 of  
rapamycin, modules 2-8 of FK-506/520 and modules 13-14 of rapamycin, wherein said  
polyketide chain is passed from module 1 of rapamycin to modules 2-8 of FK-506/520  
and then to modules 13-14 of rapamycin.

18. The polyketide synthase of claim 6 which contains modules 1-10 of  
rapamycin and modules 7-10 of FK-506/520, wherein said polyketide chain is passed  
from modules 1-10 of rapamycin to modules 7-10 of FK-506/520.

19. A method to prepare a desired polyketide which method comprises  
incubating required substrates with the polyketide synthase of claim 6.

20. The method of claim 19 wherein the substrates comprise a diketide thioester and thioesters of the required extender units.

21. The method of claim 20 wherein the extender units are malonyl, methylmalonyl, ethylmalonyl or hydroxymalonyl thioesters.

22. A polyketide or polyketide derivative synthesized by a method comprising the method of claim 19.

add  
a<sub>3</sub>

add  
a<sub>4</sub>

00000000-00000000